

The incidental capture of bats in Douglas-fir beetle funnel traps

Erin McLeod^{1,2}, Darcie Quamme³, Cori Lausen⁴, & Frances Swan^{1,2}

¹True North Forestry Consulting Ltd., ²Nakusp and Area Community Forest (NACFOR), ³Integrated Ecological Research, ⁴Wildlife Conservation Society

Project Goals

- Encourage stewardship and education on the ecological services of bats provided to the forest industry
- Collect data on incidental bat captures, providing information on bats and their prey
- Provide solutions to prevent incidental bat captures



Bats Attracted to Beetle Traps

- Long-eared bats (*Myotis evotis*) were found dead and alive
- In 2018, 4 bats were trapped in 15 weeks
- In 2019, 3 bats were trapped in 13 weeks
- Most bats were found from mid-July to mid-August
- Bats were found in traps in recent cutblocks, 50-100 m from the timber edge
- Bats are unable to get out of the trap because of the smooth surfaces and straight sides of the collection cups



Douglas-fir Beetle Funnel Traps


- Monitor and control Douglas-fir beetle (*Dendroctonus pseudotsugae*) populations in the Selkirk Resource District
- Combined effort between forest licensees and provincial government
- Pheromone-baited funnel traps collect beetles during beetle flight period (early May to late August)
- Beetles are collected in the trap, killed by insecticide, and retrieved weekly for analysis
- Traps are removed at the end of the beetle flight

Proposed Efforts to Prevent Trapping Bats

- Wire mesh placed over the bottom funnel of the trap to prevent bats from entering the collection cup
- Poles wrapped in wire mesh to act as ladders so bats can climb out of collection cups
- Enclose the whole funnel trap in wire mesh so bats can't enter the trap system but beetles will not be impeded

Data Collection and Reporting

- Multiple bat captures within a trapping grid collected over time can be uploaded to the B.C. Conservation Data Centre.
- Additional project fields within the Wildlife Incidental Observation Form should include if possible:
 - Number of pheromone traps
 - Pheromone lures used
 - Time period monitored
 - Insect trapping data
 - Location of traps with non-captures
- This would allow quantification of capture effort (% capture) and inspection of the timing of bat trapping and insect peaks



An Excel template can be obtained from SPI_Mail@gov.bc.ca

<https://a100.gov.bc.ca/pub/wiof/locationForm.do>